

cleanslate Phoenix

Disinfectant Cleaner



KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice. **IF SWALLOWED:** Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF INHALED:** Move the person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

This product is effective against SARS-CoV-2 (COVID 19 virus) on hard non-porous surfaces in 1 minute! This product is a phosphate free pH 5.5-7.5 formulation designed to provide effective cleaning, deodorizing, and disinfection¹ for hospitals², nursing homes², schools, food service establishments¹, restaurants¹, transportation terminals, office buildings, manufacturing facilities, lodging establishments, hotels, retail businesses, and athletic/recreational facilities, where housekeeping is of prime importance in reducing cross-contamination between treated surfaces. This product, when used as directed, is formulated to disinfect the following hard non-porous inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, vanity tops, shower stalls, bathtubs, and cabinets. ¹when used according to disinfection directions. ²Non-food contact use sites.

ACTIVE INGREDIENTS:

Octyl decyl dimethyl ammonium chloride.....	0.814%
Dioctyl dimethyl ammonium chloride	0.407%
Didecyl dimethyl ammonium chloride	0.407%
Alkyl (50% C14, 40% C12, 10% C16) dimethyl benzyl ammonium chloride.....	1.085%
INERT INGREDIENTS.....	97.287%
TOTAL.....	100.000%

EPA Reg. No. 1839-168-84179 EPA Est. No. 84179-NJ-1
Sold by: Cleanslate Group, LLC 1420 East Linden Ave, Linden NJ 07036

net contents: 1 Gal. (128 fl oz)



critical and semi-critical medical device surfaces. Surfaces which may contact food must be rinsed thoroughly with potable water after use. **Disinfection-** To disinfect inanimate, hard non-porous surfaces add 4 ounces of this product per gallon of water (or equivalent dilution). Apply solution with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow to remain visibly wet for 10 minutes (or 1 minute for SARS-Related Coronavirus 2 (SARS-CoV-2)), then remove excess liquid. For sprayer applications, spray 6-8 inches from surface. Do not breathe the spray mist. For visibly soiled areas, a pre-cleaning step is required. **BACTERICIDAL STABILITY OF USE-SOLUTION:** Prepare a fresh solution for each use. **To disinfect food service establishment or restaurant food contact surfaces:** floors, walls, storage areas, countertops, appliances and tables, add 4 ounces of this product per gallon of water. For visibly soiled areas, a pre-cleaning step is required. Apply solution with a cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow the surface to remain visibly wet for 10 minutes, then remove excess liquid and rinse the surfaces with potable water. This product cannot be used to clean the following food contact surfaces: utensils, glassware and dishes. **Non-Acid Toilet Bowl Disinfection/Cleaner Directions:** Remove visible filth prior to disinfection. **From Concentrate:** Add 4 ounces to the toilet bowl and mix. Use brush or sponge to apply this solution to all bowl surfaces above water line. Scrub surfaces completely, making sure to scrub under the rim, allow to stand for 10 minutes and flush. **Bactericidal Activity-** At the 4 ounce per gallon dilution, this product demonstrates effective disinfectant activity against the organisms: Pseudomonas aeruginosa PRD-10, Salmonella (choleraesuis) enterica, Staphylococcus aureus, Staphylococcus aureus (clinical isolate), Bordetella bronchiseptica, Corynebacterium ammoniagenes, Enterobacter aerogenes, Enterobacter cloacae, Enterobacter cloacae (clinical isolate), Enterococcus faecalis, Enterococcus faecalis (clinical isolate), Escherichia coli, Escherichia coli (clinical isolate), Fusobacterium necrophorum, Klebsiella pneumoniae subsp. pneumoniae, Lactobacillus casei subsp. rhamnosus, Listeria monocytogenes, Pasteurella multocida, Proteus vulgaris, Proteus mirabilis ATCC 25921, Salmonella (paratyphi B) enterica, Salmonella (typhi) enterica, Salmonella (typhimurium) enterica, Salmonella (pullorum) enterica, Serratia marcescens, Shigella sonnei, Shigella flexneri Type 2b, Shigella dysenteriae, Staphylococcus aureus subsp. aureus, Staphylococcus epidermidis, Staphylococcus epidermidis (clinical isolate), Streptococcus pyogenes (Clinical - Flesh Eating Strain BIRD M3), Streptococcus pyogenes Group A, Xanthomonas maltophilia (clinical isolate), Vancomycin resistant Enterococcus faecalis (VRE), Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin intermediate resistant Staphylococcus aureus (VISA). ***Virucidal Activity-** This product, when used on environmental, inanimate, hard, non-porous surfaces exhibits effective virucidal activity against HIV-1, HIV-2, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes Simplex Type 1 (causative agent of fever blisters), Herpes Simplex Type 2 (genital), influenza A2/Hong Kong, Vaccinia, Rotavirus, Human Coronavirus (ATCC VR-740, Strain 229E), Bovine Viral Diarrhea Virus (BVDV), Pseudorabies, Bovine Rhinotracheitis, Feline Leukemia, Feline Picornavirus, Canine Distemper Virus and SARS-Related Coronavirus 2 (SARS-CoV-2). Add 4 ounces of the product per gallon of water. For visibly soiled areas, a pre-cleaning step is required. Apply solution with a cloth, mop, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow the surface to remain visibly wet for 10 minutes (or 1 minute for SARS-Related Coronavirus 2(SARS-CoV-2)), then remove excess liquid. **Kills HIV-1, HIV-2, HBV and HCV On Pre-cleaned Environmental Surfaces/Objects Previously Soiled With Blood/body Fluids** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 or Type 2 (HIV-1 or HIV 2) (associated with AIDS), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV). **Special Instructions For Cleaning And Decontamination Against HIV-1, HIV-2, HBV and HCV Of Surfaces/Objects Soiled With Blood/Body Fluids.** **Personal Protection:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings. **Cleaning Procedures:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **Contact Time:** Allow surface to remain visibly wet for 10 minutes. **Disposal Of Infectious Materials:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal. **General Cleaning:** Apply this product to soiled area with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer. This product can be diluted at 4 ounces per gallon or applied at full strength. Thoroughly wet soiled surface and scrub as necessary. This product may be used to clean porous or non-porous surfaces. **Deodorization:** To deodorize, apply this product as indicated under the heading DISINFECTION.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. PESTICIDE STORAGE- Store in a dry place no lower in temperature than 50°F or higher than 120°F. Store in an area inaccessible to children. **PESTICIDE DISPOSAL-** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL-** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning if available. If not available, puncture and dispose in a sanitary landfill.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. KEEP OUT OF REACH OF CHILDREN. CORROSIVE. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. May be fatal if absorbed through skin. Harmful if swallowed. Wear goggles or face shield, rubber gloves, and protective clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.